

ICD-10

An Introduction

ICD-10 Worldwide

- Other countries have been using ICD-10 for many years.
- Coding professionals there were better prepared with:
 - Human Anatomy and Physiology
 - Understanding of disease process
 - Familiarity with treatments
- The above points help the coding professional better recognize and identify weaknesses in documentation required with the specificity of the new system.

Comparison

ICD-10

- 3-7 Alphanumeric
- ~68,000

ICD-9

- 3-5 Numeric (+V and E codes)
- ~13,500

Resulting in...

- An unabridged ICD-9 to ICD-10 mapping
 - Each ICD-9 code is linked to more than one ICD-10 code
 - Each ICD-10 code is more specific

General Equivalency Mapping

GEMs

Equal Mapping

ICD-10

- A02.21

Salmonella meningitis

- C92.01

Acute myeloid leukemia,
in remission

ICD-9

- 003.21

Salmonella meningitis

- 205.01

Myeloid leukemia, acute,
in remission

Unequal Mapping

ICD-10 (State of Pregnancy)

- O26.851

Spotting complicating pregnancy,
first trimester

- O26.852

Spotting complicating pregnancy,
second trimester

- O26.853

Spotting complicating pregnancy,
third trimester

ICD-9 (Episode of care)

- 649.50

Spotting complicating pregnancy,
unspecified episode of care

- 649.51

Spotting complicating pregnancy,
delivered

- 649.53

Spotting complicating pregnancy,
antepartum

One ICD-9 maps to multiple ICD-10

82002 Fracture of midcervical section of femur, closed

- From S72031A Displaced midcervical fracture of right femur, initial encounter for closed fracture
- From S72031G Displaced midcervical fracture of right femur, subsequent encounter for closed fracture with delayed healing
- From S72032A Displaced midcervical fracture of left femur, initial encounter for closed fracture
- From S72032G Displaced midcervical fracture of left femur, subsequent encounter for closed fracture with delayed healing
- *And other codes from the GEMs*

One ICD-10 maps to multiple ICD-9

- E11341 Type 2 diabetes mellitus with severe non-proliferative diabetic retinopathy with macular edema
 - 25050 Diabetes with ophthalmic manifestations, Type II or specified type, or stated as uncontrolled
 - 36206 Severe proliferative diabetic retinopathy
 - 36207 Diabetic macular edema

Some ICD-10 codes have no predecessor ICD-9 codes

- T500x6A Underdosing of mineralocorticoids and their antagonists, initial encounter
- T501x6A Underdosing of loop [high-ceiling] diuretics, initial encounter
- T502x6A Underdosing of carbonic-anhydrase inhibitors, benzothiadiazides and other diuretics, initial encounter
- T503x6A Underdosing of electrolytic, caloric and water-balance agents, initial encounter
- T504x6A Underdosing of drugs affecting uric acid metabolism, initial encounter
- *And others found in the GEMs*

Structural Differences

ICD-9-CM Diagnoses


- ICD-9-CM has 3 –5 digits
- Chapters 1 –17: all characters are numeric
- Supplemental chapters: first digit is alpha (E or V), remainder are numeric
- Decimal placed after the first three characters
 - 496 Chronic airway obstruction, not elsewhere classified (NEC)
 - 511.9 Unspecified pleural effusion
 - V02.61 Hepatitis B carrier

Structural Differences


ICD-10-CM Diagnoses

- ICD-10-CM codes have 3 –7 digits
- First digit is alpha (U is not used)
- Second and third digits are numeric
- Fourth, fifth, sixth and seventh digits may be alpha or numeric
- Decimal placed after the first three characters
 - A78 Q fever
 - A69.20 Lyme disease, unspecified
 - O9A.311 Physical abuse complicating
 pregnancy, first trimester
 - S42.001A Fracture of unspecified part of
 right clavicle, initial encounter
 for closed fracture

ICD-9-CM



ICD-10



7th Digit

- The seventh digit extensions represent
 - Visit encounter (initial = **A** or subsequent = **D**)
 - Sequelae = **S**
 - injuries and
 - external causes

7th Digits for Fractures

- **A** Initial encounter for closed fracture
- **B** Initial encounter for open fracture
- **D** Subsequent encounter for fracture with routine healing
- **G** Subsequent encounter for fracture with delayed healing
- **K** Subsequent encounter for fracture with nonunion
- **P** Subsequent encounter for fracture with malunion
- **S** Sequelae

Coding Guidelines

- Coding conventions for ICD-10-CM are found in the
 - Coding Guidelines,
 - general use guidelines, and
 - chapter-specific guidelines for the tabular list.
 - And within the index entries.
- available at
www.cdc.gov/nchs/data/icd9/draft_i10guideIn.pdf

Index

- The two parts of the ICD-10-CM index are
 - The index to diseases and injury
 - The table of drugs and chemicals and
 - The neoplasm table are housed in the index to diseases and injury.
 - The index to external causes of injury

V Codes

- V codes are now **Z codes**
- Contained in chapter 21, “Factors Influencing Health Status and Contact with Health Services.”

Improvements in Organization

- ICD-10 has 21 chapters
- Some chapters include the addition of a sixth character
- ICD-10 includes full code titles for all codes
- Z codes are now the supplemental classifications
- Post-operative complications have been moved to a procedure-specific body system chapter

NEW!

- Combination codes for conditions and common symptoms or manifestations
- Combination codes for poisonings and external causes
- Added laterality
- Added extensions for episode of care
- Expanded codes (injury, diabetes, alcohol/substance abuse, postoperative complications)

NEW!

- Inclusion of trimester in obstetrics codes and elimination of fifth digits for episode of care
- Expanded detail relevant to ambulatory and managed care encounters
- Changes in timeframes specified in certain codes
- External cause codes no longer a supplementary classification

Exclude notes defined

- Excludes1 indicates *not coded here*.
- The code being excluded is never used with the code. **The two conditions cannot occur together.**
- For example, B06 Rubella [German measles] has an Excludes1 of congenital rubella (P35.0).

Exclude notes defined

- Excludes2 indicates *not included here*.
- The excluded condition is not part of the condition represented by the code.
- It is **acceptable** to use both codes together if the patient has both conditions.
- For example, J04.0, Acute laryngitis has an Excludes2 of chronic laryngitis (J37.0).

ICD-10-CM Code Examples

- **Combination Codes for Conditions and Common Symptoms**
 - I25.110, Arteriosclerotic heart disease of native coronary artery with unstable angina pectoris
 - K50.013, Crohn's disease of small intestine with fistula
 - K71.51, Toxic liver disease with chronic active hepatitis with ascites
- **Combination Codes for Poisonings and the External Cause**
 - T39.011, Poisoning by aspirin, accidental (unintentional)
 - T39.012, Poisoning by aspirin, intentional self harm
 - T39.013, Poisoning by aspirin, assault
 - T39.014, Poisoning by aspirin, undetermined
- **Laterality**
 - C50.212, Malignant neoplasm of upper-inner quadrant of left female breast
 - H02.835, Dermatochalasis of left lower eyelid
 - I80.01, Phlebitis and thrombophlebitis of superficial vessels of right lower extremity
 - L89.213, Pressure ulcer of right hip, stage III

Diabetes Mellitus

- Includes the manifestation
 - E08.22, Diabetes mellitus due to an underlying condition with diabetic chronic kidney disease
 - E09.52, Drug or chemical induced diabetes mellitus with diabetic peripheral angiopathy with gangrene
 - E10.11, Type 1 diabetes mellitus with ketoacidosis with coma
 - E11.41, Type 2 diabetes mellitus with diabetic mononeuropathy

ICD-10 Web Resources

- General ICD-10 information

<http://www.cdc.gov/nchs/about/major/dvs/icd10des.htm>

- ICD-10-CM files, information, and General Equivalence Mappings (GEM) between ICD-10-CM and ICD-9-CM

<http://www.cdc.gov/nchs/about/otheract/icd9/icd10cm.htm>

- ICD-10-CM Official Guidelines for Coding and Reporting

http://www.cms.hhs.gov/ICD10/Downloads/7_Guidelines10cm2010.pdf

References

- Barta, Ann; et al.. "ICD-10-CM Primer." *Journal of AHIMA* 79, no.5 (May 2008): 64-66
- Centers for Medicare and Medicaid Services website

http://www.cms.hhs.gov/ICD10/01m_2009_ICD10PCS.asp